







# Whole-Genome Sequencing and *De Novo* Assembly of 61 *Staphylococcus pseudintermedius* Isolates from Healthy Dogs and Dogs with Pyoderma

 O. Francino,<sup>a</sup>  D. Pérez,<sup>a</sup>  J. Viñes,<sup>a,c</sup> R. Fonticoba,<sup>b</sup> S. Madroño,<sup>a</sup> G. Meroni,<sup>d</sup> P. Martino,<sup>d</sup> S. Martínez,<sup>e</sup>  A. Cusco,<sup>c</sup>  
 N. Fàbregas,<sup>c</sup>  L. Migura-García,<sup>f</sup> L. Ferrer<sup>b,g</sup>

<sup>a</sup>SVGM, Molecular Genetics Veterinary Service, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, Spain

<sup>b</sup>Department of Animal Medicine and Surgery, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, Spain

<sup>c</sup>Vetgenomics, Edifici EUREKA, PRUAB, Bellaterra, Barcelona, Spain

<sup>d</sup>Department of Biomedical, Surgical and Dental Sciences—One Health Unit, Milan, Italy

<sup>e</sup>Hospital Escuela de Pequeños Animales (HEPA), Facultad de Ciencias Veterinarias de la Universidad Nacional del Centro de la Provincia de Buenos Aires, Tandil, Buenos Aires, Argentina

<sup>f</sup>IRTA, Centre de Recerca en Sanitat Animal (CRESA, IRTA-UAB), Bellaterra, Barcelona, Spain

<sup>g</sup>Animal Medicine and Surgery, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, Spain

**ABSTRACT** We have *de novo* assembled and polished 61 *Staphylococcus pseudintermedius* genome sequences with Nanopore-only long reads. Completeness was 99.25%. The average genome size was 2.70 Mbp, comprising 2,506 coding sequences, 19 complete rRNAs, 56 to 59 tRNAs, and 4 noncoding RNAs (ncRNAs), as well as CRISPR arrays.

*Staphylococcus pseudintermedius* is a common microorganism of canine skin (1, 2) and the leading cause of pyoderma in dogs (3, 4). In this study, we aimed to retrieve *S. pseudintermedius* high-quality genome sequences from healthy dogs and dogs with pyoderma using a *de novo* assembly and polishing strategy with Nanopore-only long reads.

Samples were obtained by rubbing a sterile swab on the skin of healthy dogs ( $n = 22$ ; H) and the nonlesional skin of a dog with pyoderma ( $n = 6$ ; DH) or from the pustules of dogs with pyoderma ( $n = 33$ ; D). After culture in blood agar at 37°C for 24 hours, colonies were seeded in 3 ml of brain heart infusion (BHI) broth at 37°C for 16 hours. DNA was extracted with a ZymoBIOMICS DNA miniprep kit (Zymo Research, Irvine, CA, USA). DNA quality and quantity were determined using a Nanodrop 2000 spectrophotometer and Qubit double-stranded DNA (dsDNA) broad-range (BR) assay kit (Fisher Scientific SL, Madrid, Spain). The sequencing libraries were prepared with the rapid barcoding sequencing kit (SQK-RBK004; Oxford Nanopore Technologies, UK). Twelve barcoded samples were loaded into a MinION FLO-MIN106 v9.4.1 flow cell (Oxford Nanopore Technologies Ltd.) and sequenced into a MinION Mk1B instrument. The fast5 files were basecalled and demultiplexed and adapters trimmed with Guppy 4.0.11 (Oxford Nanopore Technologies) (--config dna\_r9.4.1\_450bp-s\_hac.cfg) (--config configuration.cfg --barcode\_kits SQK-RBK004 --trim\_barcode; min\_score threshold default 60). Reads with a quality score lower than 7 were discarded. Run summary statistics were obtained with Nanoplot 1.27 (5) (--N50 --fastq).

Samples assigned to *S. pseudintermedius* by WIMP (6) were *de novo* assembled using Flye 2.7.1 (7) (--nano-raw --genome-size 2.6m --plasmids --trestle). After minimap 2.17 alignment (8), the resulting contigs were polished with Racon 1.4.13 (<https://github.com/lbcb-sci/racon>) and Medaka 1.0.3 (<https://nanoporetech.github.io/medaka/>) (medaka\_consensus; -m r941\_min\_high\_g360). Genome completeness was assessed with CheckM 1.1.1 (lineage\_wf) (9). Circlator 1.5.5 was used to identify the origin (10) (fixstart --min\_id 70). Genomes were annotated with NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (11).

**Citation** Francino O, Pérez D, Viñes J, Fonticoba R, Madroño S, Meroni G, Martino P, Martínez S, Cusco A, Fàbregas N, Migura-García L, Ferrer L. 2021. Whole-genome sequencing and *de novo* assembly of 61 *Staphylococcus pseudintermedius* isolates from healthy dogs and dogs with pyoderma. Microbiol Resour Announc 10:e00152-21. <https://doi.org/10.1128/MRA.00152-21>.

**Editor** Irene L. G. Newton, Indiana University, Bloomington

**Copyright** © 2021 Francino et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to O. Francino, [olga.francino@uab.cat](mailto:olga.francino@uab.cat).

**Received** 9 February 2021

**Accepted** 22 March 2021

**Published** 22 April 2021

**TABLE 1** Characteristics and accession numbers for high-quality genome assemblies from 61 *Staphylococcus pseudintermedius* isolates<sup>a</sup>

Isolate ID <sup>b</sup>	MLST <sup>c</sup>	Source <sup>d</sup>	Yr of isolation	Country of isolation	Assembly ID (accession version no.)	Assembly level <sup>e</sup>	Genome assembly no.	Genome accession	Genome assembly	No. of contigs	N <sub>50</sub> (contigs) (bp)	Genome size (all contigs) (bp)	No. of CDS (total)	COV <sup>f</sup> (x)	Comp <sup>f</sup> (%)	Cont <sup>f</sup> (%)
DG040	Unknown	Cf, A, SP	2017	Italy	GCF_016482145.1	Contig	JAENDF0000000000		Circular	3	2,527,299	2,564,892	2,369	664	99.43	0.00
DG050	Unknown	Cf, A, SP	2016	Italy	GCF_016482445.1	Contig	JAENDE0000000000		Circular	4	2,594,061	2,637,713	2,429	326	99.43	0.57
DG059	Unknown	Cf, A, SP	2017	Italy	GCF_016481825.1	Contig	JAENDD0000000000		Circular	2	2,568,059	2,573,568	2,364	223	99.29	0.00
DG062	ST71	Cf, A, SP	2016	Italy	GCF_016482085.1	Contig	JAENDC0000000000		Linear	10	2,096,046	2,991,046	2,901	253	99.43	2.84
DG063	ST71	Cf, A, SP	2016	Italy	GCF_016482045.1	Contig	JAENDB0000000000		Linear	10	2,099,753	2,981,523	2,891	256	99.43	0.57
DG064	ST71	Cf, A, SP	2017	Italy	GCF_01645205.1	Contig	CP066718		Circular	1	2,895,060	2,895,060	2,764	361	99.43	0.57
DG066	ST71	Cf, A, SP	2017	Italy	GCF_016482585.1	Contig	JAENDA0000000000		Circular	3	2,804,246	2,808,032	2,639	213	99.43	0.57
DG067	ST71	Cf, A, SP	2017	Italy	GCF_016482535.1	Contig	JAENCZ0000000000		Linear	3	2,890,387	2,896,399	2,776	533	99.43	0.57
DG071	ST71	Cf, A, SP	2012	Italy	GCF_016482425.1	Contig	JAENCY0000000000		Linear	3	2,800,617	2,807,986	2,644	85	99.43	0.57
DG072	ST71	Cf, A, SP	2012	Italy	GCF_016455165.1	Contig	CP066717		Circular	1	2,837,133	2,837,133	2,690	304	99.43	0.57
DG076	ST258	Cf, A, SP	2012	Italy	GCF_016482525.1	Contig	JAENCX0000000000		Circular	5	2,732,273	2,780,144	2,573	323	99.43	0.85
DG077	ST71	Cf, A, SP	2016	Italy	GCF_016481985.1	Contig	JAENCW0000000000		Circular	4	2,789,419	2,791,496	2,624	479	99.43	0.57
DG078	ST71	Cf, A, SP	2012	Italy	GCF_016481935.1	Contig	JAENCV0000000000		Circular	3	2,792,180	2,799,396	2,637	208	99.43	0.57
DG081	ST301	Cf, A, SP	2012	Italy	GCF_016482005.1	Contig	JAENCU0000000000		Circular	3	2,676,449	2,694,747	2,519	353	99.43	0.00
DG082	ST258	Cf, A, SP	2017	Italy	GCF_016481945.1	Contig	JAENCT0000000000		Circular	2	2,623,528	2,626,557	2,396	259	99.43	0.00
DG089	ST71	Cf, A, SP	2012	Italy	GCF_016481925.1	Contig	JAENCS0000000000		Circular	2	2,793,862	2,797,937	2,628	281	99.43	0.57
DG093	ST258	Cf, A, SP	2017	Italy	GCF_016481905.1	Contig	JAENCR0000000000		Circular	2	2,640,063	2,648,891	2,433	39	99.43	0.00
DG094	ST71	Cf, A, SP	2017	Italy	GCF_016481855.1	Contig	JAENCQ0000000000		Circular	2	2,840,672	2,849,103	2,675	191	99.43	0.57
DG099	Unknown	Cf, A, SP	2017	Italy	GCF_016455185.1	Chrom	CP066716		Linear	1	2,623,014	2,623,014	2,416	168	99.43	0.57
DSP020	ST71	Cf, A, SP	2013	Spain	GCF_016455145.1	Contig	CP066715		Circular	1	2,793,830	2,793,830	2,637	213	99.15	0.57
DSP021	ST71	Cf, A, SP	2013	Spain	GCF_016455225.1	Contig	CP066714		Circular	1	2,795,724	2,795,724	2,637	197	99.43	0.57
DSP022	Unknown	Cf, A, SP	2014	Spain	GCF_016481865.1	Contig	JAENCP0000000000		Circular	3	2,718,542	2,767,901	2,555	174	98.86	0.28
DSP024	ST611	Cf, A, SP	2014	Spain	GCF_016482205.1	Contig	JAENCO0000000000		Circular	2	2,758,929	2,762,026	2,595	289	99.43	0.57
DSP025	ST71	Cf, A, SP	2014	Spain	GCF_016482155.1	Contig	JAENCN0000000000		Circular	6	2,801,923	2,805,515	2,641	295	99.43	0.57
DSP026	ST503	Cf, A, SP	2014	Spain	GCF_016455085.1	Contig	CP066713		Circular	1	2,567,628	2,567,628	2,387	303	99.43	0.00
DSP027	Unknown	Cf, A, SP	2014	Spain	GCA_016455285.1	Contig	CP066712		Circular	1	2,717,194	2,717,194	2,537	352	99.43	0.28
DSP028	Unknown	Cf, A, SP	2019	Spain	GCF_016482125.1	Contig	JAENCM0000000000		Circular	2	2,569,147	2,575,420	2,367	238	98.86	0.00
DSP029	ST258	Cf, A, SP	2019	Spain	GCF_016481265.1	Contig	JAENBQ0000000000		Linear	8	2,413,604	2,723,805	2,517	270	99.43	0.00
DSP030	Unknown	Cf, A, SP	2018	Argentina	GCF_016455265.1	Contig	CP066711		Circular	1	2,612,059	2,612,059	2,386	164	99.43	0.00
DSP032	ST1631	Cf, A, SP	2019	Argentina	GCA_016482035.1	Contig	JAENCL0000000000		Linear	3	1,369,210	2,670,199	2,449	119	98.66	0.57
DSP034	ST1827	Cf, A, SP	2018	Argentina	GCF_016455025.1	Contig	CP066710		Circular	1	2,550,368	2,550,368	2,319	204	99.43	0.00
DSP035	Unknown	Cf, A, SP	2019	Argentina	GCF_016481845.1	Contig	JAENCK0000000000		Linear	2	2,640,488	2,642,865	2,489	117	99.43	0.00
DSP036	ST71	Cf, A, SP	2019	Spain	GCF_016482105.1	Contig	JAENCJ0000000000		Circular	7	2,849,760	2,927,015	2,755	129	99.43	0.57
DHSP041	ST71	Cf, A, H	2019	Spain	GCF_016482265.1	Contig	JAENBW0000000000		Linear	3	2,849,717	2,909,751	2,748	123	99.41	0.57
DHSP042	ST71	Cf, A, H	2019	Spain	GCF_016482345.1	Contig	JAENBV0000000000		Linear	6	2,822,064	2,930,017	2,768	72	99.43	0.57
DHSP043	ST71	Cf, A, H	2019	Spain	GCF_016482305.1	Contig	JAENBU0000000000		Linear	12	1,649,358	2,931,289	2,776	66	99.43	0.57
DHSP044	ST71	Cf, A, H	2019	Spain	GCF_016482225.1	Contig	JAENBT0000000000		Circular	4	2,849,602	2,907,542	2,748	132	99.43	0.57
DHSP045	ST71	Cf, A, H	2019	Spain	GCF_016482025.1	Contig	JAENBS0000000000		Circular	5	2,849,736	2,915,461	2,742	127	99.41	0.57
DHSP046	ST496	Cf, P, H	2019	Spain	GCF_016481295.1	Contig	JAENBR0000000000		Circular	2	2,695,721	2,710,903	2,475	100	99.43	0.00
HSP079	Unknown	Cf, P, H	2019	Spain	GCF_016598895.1	Contig	CP066885		Circular	1	2,585,691	2,587,693	2,337	301	99.43	0.00
HSP080	Unknown	Cf, P, H	2019	Spain	GCF_016598915.1	Contig	CP066884		Circular	1	2,585,570	2,587,506	2,336	199	99.41	0.00
HSP081	Unknown	Cf, P, H	2019	Spain	GCA_016482135.1	Contig	JAENCI0000000000		Circular	4	2,617,897	2,621,254	2,467	94	98.58	1.14

(Continued on next page)

TABLE 1 (Continued)

Isolate ID <sup>b</sup>	MLST <sup>c</sup>	Source <sup>d</sup>	Yr of isolation	Country of isolation	Assembly ID (accession version no.)	Assembly level <sup>e</sup>	Genome accession no.	Genome assembly	No. of contigs	N <sub>50</sub> (contigs) (bp)	Genome size (all contigs) (bp)	No. of CDS (total)	COV <sup>f</sup> (x)	Comp <sup>f</sup> (%)	Cont <sup>f</sup> (%)
HSP082	Unknown	Cf, P, H	2019	Spain	GCA_016481725.1	Contig	JAENCH0000000000	Circular	2	2,617,857	2,620,663	2,466	149	96.73	1.14
HSP093	Unknown	Cf, A, H	2019	Spain	GCF_016481735.1	Contig	JAENCG0000000000	Circular	3	2,571,836	2,590,335	2,361	274	99.43	0.57
HSP094	Unknown	Cf, A, H	2019	Spain	GCF_016481805.1	Contig	JAENCF0000000000	Circular	2	2,567,494	2,570,595	2,337	550	99.43	0.57
HSP095	Unknown	Cf, A, H	2019	Spain	GCF_016482325.1	Contig	JAENCE0000000000	Circular	2	2,570,328	2,575,879	2,360	341	99.43	0.57
HSP096	Unknown	Cf, A, H	2019	Spain	GCF_016482245.1	Contig	JAENCD0000000000	Circular	3	2,570,206	2,578,330	2,359	89	99.43	0.57
HSP097	Unknown	Cf, A, H	2019	Spain	GCF_016481665.1	Contig	JAENCC0000000000	Circular	3	2,570,223	2,575,223	2,356	136	99.43	0.57
HSP118	Unknown	Cf, A, H	2019	Spain	GCF_016455005.1	Compl	CP066709	Circular	1	2,512,855	2,512,855	2,277	313	99.43	0.00
HSP125	ST1248	Cf, A, H	2019	Spain	GCF_016455245.1	Compl	CP066708	Circular	1	2,551,473	2,551,473	2,330	327	98.86	0.00
HSP127	ST1061	Cf, P, H	2019	Spain	GCF_016481685.1	Contig	JAENCB0000000000	Circular	2	2,660,509	2,690,618	2,507	162	98.86	1.14
HSP132	Unknown	Cf, P, H	2019	Spain	GCA_016455125.1	Chrom	CP066707	Linear	1	2,515,164	2,515,164	2,275	132	98.72	0.00
HSP134	Unknown	Cf, P, H	2019	Spain	GCA_016455105.1	Compl	CP066706	Circular	1	2,514,594	2,514,594	2,274	146	98.86	0.00
HSP135	Unknown	Cf, A, H	2019	Spain	GCF_016455065.1	Compl	CP066705	Circular	1	2,512,727	2,512,727	2,277	154	98.86	0.00
HSP136	Unknown	Cf, A, H	2019	Spain	GCA_016455045.1	Chrom	CP066704	Linear	1	2,512,726	2,512,726	2,274	104	98.86	0.00
HSP137	Unknown	Cf, A, H	2019	Spain	GCF_016454985.1	Compl	CP066703	Circular	1	2,512,757	2,512,757	2,271	224	99.43	0.00
HSP138	Unknown	Cf, A, H	2019	Spain	GCF_016454965.1	Compl	CP066702	Circular	1	2,512,830	2,512,830	2,277	227	98.86	0.00
HSP140	Unknown	Cf, P, H	2019	Spain	GCF_016482365.1	Contig	JAENCA0000000000	Circular	2	2,594,004	2,597,272	2,387	296	99.24	0.00
HSP141	ST257	Cf, P, H	2019	Spain	GCF_016482355.1	Contig	JAENBZ0000000000	Circular	4	2,615,633	2,622,529	2,450	150	99.43	0.57
HSP142	Unknown	Cf, P, H	2019	Spain	GCF_016482405.1	Contig	JAENBY0000000000	Circular	2	2,594,059	2,597,098	2,393	461	99.43	0.00
HSP143	Unknown	Cf, P, H	2019	Spain	GCF_016482235.1	Contig	JAENBX0000000000	Linear	6	2,537,079	2,779,670	2,589	293	98.86	0.57

<sup>a</sup> Obtained from the lesional skin of 33 dogs with pyoderma (33 D), the nonlesional skin of a dog with pyoderma (6 DH), and 6 healthy dogs (22 H).<sup>b</sup> ID, identifier.<sup>c</sup> MLST, multilocus sequence type; ST, sequence type.<sup>d</sup> Cf, *Canis lupus familiaris*; A, abdominal skin swab; P, perioral skin swab; SP, superficial pyoderma; H, healthy.<sup>e</sup> Compl, complete genome; Chrom, chromosome.<sup>f</sup> COV, coverage; Comp, completeness; Cont, contamination.

Multilocus sequence types (MLSTs) were assigned with software MLST 2.0. and database 2.0.0 (<https://cge.cbs.dtu.dk/services/MLST-2.0/>) (12).

Nanopore sequencing allowed successful *de novo* assembly and polishing of 61 *S. pseudintermedius* isolates (Table 1). The average read  $N_{50}$  value was 4,223.38 bp for 2,848,339.50 reads per flow cell (237,361.63 per barcoded sample). The mean genome coverage was  $235\times$  ( $39\times$  to  $664\times$ ), with an average contig  $N_{50}$  value of 2.6 Mbp (1.6 Mbp to 2.9 Mbp). The average genome completeness was 99.25% (98.6% to 99.4%, except for HSP082), which is close to previous results with hybrid assemblies (13). The number of contigs per isolate ranged from 1 to 12 (median, 2). The main contig was circular for 47 isolates. The average genome size of *S. pseudintermedius* was 2.70 Mbp (2.51 to 2.99 Mb), comprising 2,506 coding sequences (CDS; 2,271 to 2,901), 19 complete rRNAs (6 to 7 5S, 6 16S, and 6 23S rRNA genes), 56 to 59 tRNAs, and 4 noncoding RNAs (ncRNAs), as well as CRISPR arrays (0.5; range from 0 to 2). Pangenome analyses of isolates from healthy and diseased individuals will help unravel the differences, if any, that exist between commensal and pathogenic *S. pseudintermedius* populations.

**Data availability.** The standardized strain descriptions and accession numbers are presented in Table 1; the genome sequence assemblies and genomic data are publicly available in DDBJ/ENA/GenBank under BioProject no. [PRJNA685966](#) with the accession numbers [CP066702](#) to [CP066718](#), [CP066884](#), [CP066885](#), and [JAENBQ000000000](#) to [JAENDF000000000](#). The versions described in this paper are the first versions. The raw data are available from the Sequence Read Archive (SRA) under the same BioProject no., [PRJNA685966](#).

## ACKNOWLEDGMENTS

Funding sources were as follows: Research Project RTI2018-101991-B-100 ("From whole genome sequencing to clinical metagenomics: investigations on the pathogenesis of *Staphylococcus pseudintermedius* pyoderma in the dog"); Spanish Ministry of Science, Innovation and Universities; Industrial Doctorate Program grant 2017DI037 AGAUR; and Generalitat de Catalunya, Spain. The Spanish Ministry of Science and Innovation granted a Torres Quevedo Project to Vegenomics, S.L. with reference PTQ2018-009961 that is cofinanced by the European Social Fund.

L.F. has received unrelated honoraria for lecturing from Zoetis, Bayer, LETI, and Affinity Petcare. O.F. and A.C. have received unrelated honoraria for lecturing from Oxford Nanopore Technologies.

## REFERENCES

- Paul NC, Bärman SC, Moodley A, Nielsen SS, Guardabassi L. 2012. *Staphylococcus pseudintermedius* colonization patterns and strain diversity in healthy dogs: a cross-sectional and longitudinal study. *Vet Microbiol* 160:420–427. <https://doi.org/10.1016/j.vetmic.2012.06.012>.
- Rubin JE, Chirino-Trejo M. 2011. Prevalence, sites of colonization, antimicrobial resistance among *Staphylococcus pseudintermedius* isolated from healthy dogs in Saskatoon, Canada. *J Vet Diagn Invest* 23:351–354. <https://doi.org/10.1177/104063871102300227>.
- Bannoehr J, Guardabassi L. 2012. *Staphylococcus pseudintermedius* in the dog: taxonomy, diagnostics, ecology, epidemiology and pathogenicity. *Vet Dermatol* 23:253–266, e51–2. <https://doi.org/10.1111/j.1365-3164.2012.01046.x>.
- Loeffler A, Lloyd DH. 2018. What has changed in canine pyoderma? A narrative review. *Vet J* 235:73–82. <https://doi.org/10.1016/j.tvjl.2018.04.002>.
- De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. 2018. NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* 34:2666–2669. <https://doi.org/10.1093/bioinformatics/bty149>.
- Juul S, Izquierdo F, Hurst A, Dai X, Wright A, Kulesha E, Pettett R, Turner DJ. 2015. What's in my pot? Real-time species identification on the MinION. *bioRxiv* <https://doi.org/10.1101/030742>.
- Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs. *Nat Biotechnol* 37:540–549. <https://doi.org/10.1038/s41587-019-0072-8>.
- Li H. 2018. minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100. <https://doi.org/10.1093/bioinformatics/bty191>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Hunt M, De Silva N, Otto TD, Parkhill J, Keane J, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Pontén T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>.
- Viñes J, Cuscó A, Francino O. 2020. Hybrid assembly from a pathogenic methicillin- and multidrug-resistant *Staphylococcus pseudintermedius* strain isolated from a case of canine otitis in Spain. *Microbiol Resour Announc* 9:e01121-19. <https://doi.org/10.1128/MRA.01121-19>.